

Oligoclonal Bands IgG

Order Name: **OLIGO CSF**

Test Number: 0804040

Revision Date: 12/12/2022

TEST NAME	METHODOLOGY	LOINC CODE
Oligoclonal Bands, CSF	Isoelectric Focusing	12782-9

SPECIMEN REQUIREMENTS				
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	0.5 mL (0.1 mL) Serum and CSF ea.	CSF and Serum	Sterile Screwtop Container and Clot Activator SST	Room Temperature
Instructions	Notes: Collect BOTH 0.5 mL (0.1 mL) CSF AND 0.5 mL (0.1 mL) Serum Specimen Type: Red-top tube or gel-barrier tube AND plastic (CSF) tube Specimen Storage: Room temperature Specimen Collection: Spinal tap AND venipuncture. Specimen Stability: Ambient: 14 days, Refrigerated : 14 days, Frozen: 6 months			

GENERAL INFORMATION	
Expected TAT	4-6 Days
Clinical Use	Oligoclonal bands are present in the CSF of more than 85% of patients with clinically definite multiple sclerosis (MS). To distinguish between oligoclonal bands in the CSF due to a peripheral gammopathy and oligoclonal bands due to local production in the CNS, serum and CSF should be tested simultaneously. Oligoclonal bands can however be observed in a variety of other diseases, e.g., subacute sclerosing panencephalitis, inflammatory polyneuropathy, CNS lupus, and brain tumors and infarctions.
Performing Labcorp Test Code	019216
CPT Code(s)	83916
Lab Section	Reference Lab