Striated Muscle Antibody Screen with Titer

Order Name: STRIAT AB
Test Number: 3805400
Revision Date: 02/10/2023

TEST NAME			METHODOLOGY	LOINC CODE
Striated Muscle Antibody Screen			Semi-Quantitative Indirect Fluorescent Antibody	5372-8
SPECIMEN REQUIR	EMENTS			
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	1 mL (0.5 mL)	Serum	Clot Activator SST	Refrigerated
Alternate 1	1 mL (0.5 mL)	Serum	Clot Activator (Red Top, No-Gel)	Refrigerated
Instructions	Minimum Volume: 0.5 mL (Note: This volume does not allow for repeat testing. Specimen Type: Gel-barrier tube or Red-top tube Specimen Storage: Refrigerated Specimen Collection: Separate serum from cells ASAP or within 2 hours of collection Specimen Stability: Ambient: 14 days, Refrigerated: 14 days, Frozen: 14 days (Freeze/thaw cycles Stable x3) Causes for Rejection: Hemolyzed or lipemic serum			
	If being collected for Lambert-Eaton Syndrome Panel, Please collect Serum from Red No-Gel Clot Tube.			

GENERAL INFORMATION		
Expected TAT	2 - 4 days	
Clinical Use	This test is recommended for differential evaluation of neuromuscular junction diseases including myasthenia gravis (MG). This test will report a Negative Screen and if Positive will report out the Titer. Reference Interval Negative: <1:100	
Performing Labcorp Test Code	160184	
Notes	Labcorp Test Code: 160184	
CPT Code(s)	86255	
Lab Section	Reference Lab	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025