Neisseria gonorrhoeae Urine - TMA

Order Name: GC Urine
Test Number: 6910117
Revision Date: 07/11/2017

TEST NAME			METHODOLOGY	LOINC CODE	
Neisseria gonorrhoeae Urine			Transcription-Mediated Amplification	60256-5	
SPECIMEN REQUIREMENTS					
Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment	
Preferred	2 mL	Urine, First Void	APTIMA Urine Transport Tube	Refrigerated	
Instructions	First catch urine. Patient should not have urinated for at least 1 hr prior to collection. Female pateints should not cleanse the labial area prior to collection. Transfer 2 mL urine to APTIMA Urine Specimen Transport Tube within 24hrs of collection. Liquid level must be between fill lines on tube. Stability: Refrigerated = 30 days.				
	Please contact your account	t rep if training and APTIMA U	rine Specimen Transport supplies have not be	een supplied to your practice.	

GENERAL INFORMATION		
Testing Schedule	Mon-Fri	
Expected TAT	1-2 days after set-up	
CPT Code(s)	87591	
Lab Section	Molecular Diagnostic	

Service provided by Labcorp Oklahoma, Inc. All Rights Reserved. © 2003 - 2025