

## Mycoplasma genitalium, NAA, Urine

Order Name: **M genital, UR**  
Test Number: 5195311  
Revision Date: 09/12/2023

TEST NAME	METHODOLOGY	LOINC CODE
Mycoplasma genitalium, NAA, Urine	Nucleic acid amplification (NAA)	88226-6

### SPECIMEN REQUIREMENTS

Specimen	Specimen Volume (min)	Specimen Type	Specimen Container	Transport Environment
Preferred	<b>30 mL (2 mL)</b>	<b>Urine (See Inst.)</b>	<b>APTIMA Urine Transport Tube</b>	<b>Room Temperature</b>
Instructions	<p><b>Specimen:</b> First-void urine in Aptima® urine transport</p> <p><b>Volume:</b> 2 mL of a 20 to 30 mL urine collection</p> <p><b>Container:</b> Aptima® urine transport</p> <p><b>Collection:</b> Urine specimen: The patient should not have urinated for at least one hour prior to specimen collection. Direct patient to provide a first-catch urine (approximately 20 to 30 mL of the initial urine stream) into a urine collection cup free of any preservatives. Collection of larger volumes of urine may result in specimen dilution that may reduce test sensitivity; lesser volumes may not adequately rinse organisms into the specimen. Female patients should not cleanse the labial area prior to providing the specimen. Add urine to the Aptima® Combo 2 urine collection device. The final volume must be between the two black lines on the device (about 2 mL).</p> <p><b>Storage Instructions:</b> Maintain specimen at room temperature or refrigerate (2°C to 30°C).</p> <p><b>Stability:</b> Room temperature 30 days (stability provided by manufacturer or literature reference)</p>			

### GENERAL INFORMATION

Expected TAT	3-4 days
Performing Labcorp Test Code	180025
Notes	
RESULT CODE	RESULT NAME
6907667	M genitalia Ur
CPT Code(s)	87563
Lab Section	Reference Lab